



# Las Vegas, Nevada – 2002 (N=52)


**Figure A. Age of GISP participants, in years, 2002**




**Figure B. Race/ethnicity of GISP participants, 2002**



**Figure C. Percentage of GISP participants identifying as men who have sex with men, 1988–2002**




**Figure D. Reason for visit among GISP participants, 2002**



# Las Vegas, Nevada – 2002 (N=52)

**Figure E. Previous episode of gonorrhea among GISP participants, 1991 – 2002**




\*Data first collected in 1991.


\*\*Data first collected in 1992.

Note: Data points not shown when >30% data missing.


**Figure F. Drugs used to treat gonorrhea among GISP participants, 2002**



**Figure G. Drugs used to treat *Chlamydia trachomatis* infection among GISP participants, 2002**



**Figure H. Resistance to penicillin and tetracycline among GISP isolates, 2002**



## Las Vegas, Nevada – 2002 (N=52)

**Figure I. Decreased susceptibility to ceftriaxone among GISP isolates, 1988 – 2002**

No isolates with decreased susceptibility to ceftriaxone have been identified at this clinic.

**Figure J. Decreased susceptibility to cefixime among GISP isolates, 1992 – 2002**

No isolates with decreased susceptibility to cefixime have been identified at this clinic.

**Figure K. Intermediate resistance and resistance to ciprofloxacin among GISP isolates, 1990 – 2002**

No isolates with intermediate resistance or resistance to ciprofloxacin have been identified at this clinic.

**Figure L. Decreased susceptibility to azithromycin among GISP isolates, 1992 – 2002**

No isolates with decreased susceptibility to azithromycin have been identified at this clinic.